

Microsecond simulations of the folding/unfolding thermodynamics of the Trp-cage miniprotein

Ryan Day, Dietmar Paschek, and Angel E. Garcia*

Department of Physics, Applied Physics and Astronomy and Center for Biotechnology and Interdisciplinary Studies, Rensselaer Polytechnic Institute, Troy, New York

ABSTRACT

We study the unbiased folding/unfolding thermodynamics of the Trp-cage miniprotein using detailed molecular dynamics simulations of an all-atom model of the protein in explicit solvent using the Amberff99SB force field. Replica-exchange molecular dynamics simulations are used to sample the protein ensembles over a broad range of temperatures covering the folded and unfolded states at two densities. The obtained ensembles are shown to reach equilibrium in the 1 μ s/replica timescale. The total simulation time used in the calculations exceeds 100 μ s. Ensemble averages of the fraction folded, pressure, and energy differences between the folded and unfolded states as a function of temperature are used to model the free energy of the folding transition, $\Delta G(P, T)$, over the whole region of temperatures and pressures sampled in the simulations. The $\Delta G(P, T)$ diagram describes an ellipse over the range of temperatures and pressures sampled, predicting that the system can undergo pressure-induced unfolding and cold denaturation at low temperatures and high pressures, and unfolding at low pressures and high temperatures. The calculated free energy function exhibits remarkably good agreement with the experimental folding transition temperature ($T_f = 321$ K), free energy, and specific heat changes. However, changes in enthalpy and entropy are significantly different than the experimental values. We speculate that these differences may be due to the simplicity of the semiempirical force field used in the simulations and that more elaborate force fields may be required to describe appropriately the thermodynamics of proteins.

Proteins 2010; 78:1889–1899.
© 2010 Wiley-Liss, Inc.

Key words: pressure denaturation; cold denaturation; folding; molecular simulations; force field.

INTRODUCTION

The Trp-cage miniprotein is a designed peptide that behaves like larger globular proteins.¹ This protein is stabilized by a hydrophobic core containing a Trp amino acid, whose hydrogen bonds to a distant backbone carbonyl, and by an ion pair. Structurally, the Trp-cage is composed of an alpha helix, a 3–10 helix, and a polyproline II segment. The folding is cooperative and occurs in the microsecond timescale.² The Trp-cage miniprotein has also served as an ideal model in computational studies where force fields^{3–5} and modeling techniques^{6–8} have been benchmarked against detailed structural,¹ thermodynamics,⁹ kinetics,^{2,10–12} and design^{13–16} data. So far multiple studies describing the folding of Trp-cage using atomic force fields with implicit and explicit solvent^{3,6–8,17–25} have successfully reproduced the structure of the folded state with excellent accuracy. Here, we report the equilibrium thermodynamics of the Trp-cage miniprotein that have been calculated with the recently developed Amberff99SB (ff99SB) force field⁴ and compared with thermodynamic studies.⁹ We find that the ff99SB force field and the TIP3P water model accurately describe the thermodynamics of the system and predict a folding transition at 48°C (321 K) with the specific heat of the transition in agreement with the experimental value. These results contrast with extreme stability and high folding temperature ($T_f \sim 450$ K) predicted by the Cornell *et al.* force field (ff94)²⁶ and obtained by us in similar calculations.^{18,24} We show that the ff99SB significantly improves the description of folding thermodynamics of the protein. However, the enthalpy and entropy changes during the folding/unfolding transition are underestimated by at least a factor of two. Here, we characterize the folding thermodynamics and structural ensembles of the protein as a function of temperature.

METHODS

We used replica-exchange molecular dynamics (REMD) to model the configurational ensemble of a system composed of the 20 amino

Additional Supporting Information may be found in the online version of this article.
Grant sponsor: National Science Foundation; Grant number: MCB-0543769

Ryan Day's current address is Department of Chemistry, University of the Pacific, Stockton, California

*Correspondence to: Angel E. Garcia, 1C25JR Science Center, Department of Physics, Rensselaer Polytechnic Institute, Troy, NY 12180. E-mail: angel@rpi.edu

Received 7 October 2009; Revised 6 January 2010; Accepted 17 January 2010

Published online 16 February 2010 in Wiley InterScience (www.interscience.wiley.com).

DOI: 10.1002/prot.22702

acid Trp-cage (Ac-NLYIQWLKDGPPSSGRPPPS-Nme) with charged Lys, Arg, and Asp sidechains, one Na^+ , two Cl^- ions to neutralize the system, and 2635 TIP3P water molecules.²⁷ Electrostatics interactions were modeled using PME using a cubic $36 \times 36 \times 36$ grid and Van der Waals interactions were truncated at 1.0 nm. A grid size of 0.12 nm was used for the PME. Following the results of Cooke and Schmidler regarding the ergodicity of the sampling in REMD,²⁸ the equations of motion were integrated using the stochastic dynamics with coupling time of 1.0 ps and using a 2-fs time step. Simulations are done using GROMACS and the ff99SB force field.⁴ REMD simulations are done at constant volume in a cubic box of 4.42 nm corresponding to the volume of the system at pressure, $P = 1$ atm, and temperature, $T = 300$ K, obtained from a 10 ns NPT simulation. We simulated 40 systems with temperatures 280.0, 284.1, 288.2, 292.4, 296.7, 301.1, 305.6, 310.2, 314.9, 319.7, 324.6, 329.6, 334.7, 340.0, 345.4, 351.0, 356.6, 362.5, 368.4, 374.6, 380.9, 387.3, 394.0, 400.8, 407.8, 415.1, 422.5, 430.1, 438.0, 446.0, 454.3, 462.8, 471.6, 480.6, 489.8, 499.3, 509.0, 519.0, 529.2, and 539.7 K. Temperatures were selected such that an exchange rate of 0.15 is obtained.²⁹ REMD simulations were started from random unfolded states [we label this as “unfolded initial conditions” (UIC)] and from the folded state of the protein [we label this as “folded initial conditions” (FIC)]. The UIC states were generated by starting from an extended all-PPII conformation, which gets slightly compacted after running a short (25 ps) simulation in the gas phase at 300 K, as described in the previous calculations.²⁴ The resulting systems were solvated with a pre-equilibrated box of TIP3P water molecules at 300 K. Simulations were extended for 1 μs /replica for UIC and 0.5 μs for FIC. The length of the simulations is dictated by the relaxation behavior (of ~ 200 ns) of the system. Previous ff94 simulations were extended to 200 ns, since they showed fast (30 ns) relaxation time.²⁴ Configurations were collected for further analysis for every 1 ps. Analysis presented here use the last 0.5 and 0.25 μs for the UIC and FIC simulations, respectively. To fit the thermodynamics of the Trp-cage over a broad range of T and P , we also simulated the system at high average pressure over the same temperatures (in a cubic box of 4.176 nm). The initial configuration for this simulation was the folded state (since it reaches steady state faster than simulation started from the unfolded state), but the volume was reduced such that the average pressure at 310 K is 400 MPa. The initial state of the system was obtained from a 10 ns NPT simulation. The REMD simulation was extended for 1 μs /replica and the last 0.5 μs are used for calculating averages.

To obtain a pressure–temperature stability diagram, we fit the calculated fraction folded, the average pressure, the average potential energy, and the free energy differences as a function of temperature. Here, we combine the

averages calculated for the protein under different conditions. The free energy difference between the folded and unfolded state as a function of P and T can be written as

$$\begin{aligned} \Delta G(T, P) = & \Delta G_U - \Delta S_U(T - T_0) \\ & - \Delta C_p \left[T \left(\text{Log} \left(\frac{T}{T_0} \right) - 1 \right) + T_0 \right] + \Delta V_U(P - P_0) \\ & + \frac{1}{2} \Delta \beta_U(P - P_0)^2 + \Delta \alpha_U(P - P_0)(T - T_0) \quad (1) \end{aligned}$$

where ΔC_p is the heat capacity, ΔS_U is the entropy, ΔG_U is the Gibbs free energy, $\Delta \beta_U$ is the compressibility change, ΔV_U is the volume change, and $\Delta \alpha_U$ is the expansivity change on unfolding of the protein.³⁰ The coefficients are calculated at a reference T_0 (298 K) and P_0 (0.1 MPa = 1 atm). Each one of the six coefficients in this expansion is a measurable thermodynamic quantity. The system free energy difference is fitted by minimizing a χ^2 function as a function of the expansion coefficients. The χ^2 function is given by

$$\begin{aligned} \chi^2 = & \sum_i \left[\frac{\Delta G_i - \Delta G(T_i, P_i)}{\sigma_{\Delta Y_i}} \right]^2 \\ & + \sum_i \left[\frac{\Delta x_i^{\text{folded}} - \Delta x(T_i, P_i)}{\sigma_{\Delta x_i}} \right]^2 + \sum_i \left[\frac{\Delta E_i - \Delta E(T_i, P_i)}{\sigma_{\Delta E_i}} \right]^2 \quad (2) \end{aligned}$$


Here, $\sigma_{\Delta Y_i}$ is the uncertainty in the average of Y (where Y is the energy, free energy or fraction folded), x_i^{folded} is the fraction of molecules folded at the state i , and ΔG_i , ΔE_i are the free energy and energy changes, respectively, calculated from ensemble averages. ΔE_i is the average potential energy. The index i labels the replica states. Each replica state samples a temperature, T_i , at constant system volume. $\Delta G(T_i, P_i)$, $\Delta V(T_i, P_i)$, and $\Delta E(T_i, P_i)$ are the fitted functions evaluated for each of the replica states, (T_i, P_i) , and P_i is the average pressure. The volume change is defined as the partial derivative of the Gibbs free energy with respect to P (at constant temperature),

$$\begin{aligned} \Delta V(P, T) = & \left(\frac{\partial \Delta G(P, T)}{\partial P} \right)_T = \Delta V_U + \Delta \alpha_U(T - T_0) \\ & + \Delta \beta_U(P - P_0) \quad (3) \end{aligned}$$

The enthalpy is given by $\Delta H_i = \Delta E_i + P_i \Delta V_i$. The average energy is given by

$$\begin{aligned} \Delta E(T, P) = & \Delta H - P \Delta V = \Delta G(T, P) \\ & + T \Delta S(T, P) - P \Delta V(T, P) \quad (4) \end{aligned}$$

where


Figure 1

Time evolution of the REMD trajectories. (A) Fraction of the time that each replica (1–40) spends in the folded state ($rmsd < 0.22$ nm) over 1 ns time block averages. All replicas were started from an unfolded configuration. (B) First folding time for replicas along the folding trajectory (red line). The profile shows an exponential growth with exponent of the order of 100 ns. The green line shows the first time a replica has folded, unfolded past 0.6 nm, and refolded. The curve is very similar to the first folding time, except for a time delay of tens of nanoseconds. (C) Fraction of replicas that are folded at any time during the simulation as a function of time [red line (UIC) and green line (FIC)]. All replicas at different temperatures are averaged together. We also show the fraction of replicas that are folded for a simulation in which all replicas were started in the folded state. The two ensembles show similar averages after 200 ns. The blue curve in the inset of (C) is the time correlation function of the fraction folded time history. Estimates of the correlation time for the fraction for replicas folded are 50 ns. We use simulation segments if twice this time, 100 ns, to estimate uncertainties in the averages.

$$\Delta S(T, P) = - \left(\frac{\partial \Delta G(T, P)}{\partial T} \right)_P. \quad (5)$$

Then, the energy difference is given by

$$\begin{aligned} \Delta E(T, P) = & \Delta G_U + T_0 \Delta S_U - \Delta V_U P_0 + \Delta C_p (T - T_0) \\ & - \Delta \alpha_U (PT - P_0 T_0) - \Delta \beta_U (P^2 - P_0^2) \end{aligned} \quad (6)$$

The free energy, $\Delta G(P, T)$, is calculated from the fraction of folded states, x_i^{folded} , obtained from the sampling and is related to the free energy of folding by

$$\Delta G_i(T_i, P_i) = -RT \text{Log} \left[\frac{1 - x_i^{\text{folded}}}{x_i^{\text{folded}}} \right] \quad (7)$$

Here, R is the gas constant. Uncertainties in fraction folded, free energy, and energy are calculated from five,

100 ns segments for low and high average pressure simulations. The length of the segments, 100 ns, are taken to be about twice the correlation time of the number of folded replicas as a function of time. Fittings are done in Mathematica using steepest descent and Newton–Raphson minimization algorithms.

We estimate the correlation time, τ_c , of the number of folded replicas from the time correlation functions, $c(t)$. We obtained two estimates of the correlation time. One estimate was obtained by integrating the correlation function over time, $\tau_c = \text{Max}[\int_0^t dt' c(t')]$, until a maximum is reached and taking this maximum to be an upper bound for the correlation time. Another estimate was obtained by assuming that the errors are Gaussian distributed and calculating the integral $\tau_c = \int_0^\infty dt' c^2(t')^{31}$. Both estimates give $\tau_c \sim 50$ ns. We use segments of twice that time, 100 ns, to estimate uncertainties in the averages. We found that even 1 μ s




Figure 2

Temperature dependence of distributions of (a) $rmsd$, (b) Trp-N ϵ to Asp backbone carbonyl hydrogen bond, and (c) Arg-Asp ion pair distances. All distances are in nanometers.

simulation is not long enough to obtain better estimates of the correlation of a slowly varying function, like the number of folded replicas as a function of time.

RESULTS

Steady state equilibrium of the system

First we establish that the ensemble averages have reached a steady state by monitoring the rate at which

replicas reach the folded state, unfold and refold [Fig. 1(b)], the average number of replicas that are folded as a function of simulation time [Fig. 1(c)], and the time history of the folding probability of each replica [Fig. 1(a)]. We distinguish folded from unfolded based on the $rmsd$ distance of the configuration from the folded state reported in the PDB [NMR structure #1 (PDB-code 1L2Y)]. A configuration is considered to be folded if $rmsd < 0.22$ nm. We will justify this choice later in the manuscript using the distribution of $rmsd$ at various temperatures. To identify possible dependence of the results on initial conditions, we performed two simulations starting from the folded initial configurations (FIC) and unfolded initial configurations (UIC). For simulations with UIC all replicas have folded after $0.60 \mu\text{s}$. The refolding curve monitors the rate at which replicas refold after leaving the folded basin and reaching an $rmsd > 0.6$ nm from the folded state. This curve shows a short delay from the first folded time curve, illustrating that the averages obtained in long timescale simulations are not sensitive to initial conditions. Figure 1(c) shows that the number of folded replicas in the UIC and FIC converge after $0.20 \mu\text{s}$. The number of folded replicas shows a long correlation time of 50 ns. Similar calculations for ff94 showed that all replicas folded after 30 ns, which is an order of magnitude faster than for the ff99SB calculations presented here, and, therefore, require that the ff99SB calculations be extended to the μs timescale. The time evolution of the fraction of time, within 1 ns time blocks, that a given replica samples the folded state, shown in Figure 1(a), shows that replicas remain folded for hundreds of nanoseconds, which explain the slow relaxation time of the total number of folded replicas. Supporting Information Figure S8 shows a figure similar to Figure 1 for the high density REMD simulations.

Structural characterization of the ensembles

The structural ensembles of the protein are characterized using different order parameters. In general, not all order parameters are equally suited to describe the folding/unfolding equilibrium. In particular, the folding thermodynamics of proteins is not strictly two state and different order parameters will monitor changes differently, as is the case when using experimental probes. Ideally, an order parameter should be related to a measurable quantity like the fluorescence quenching or the strength of a circular dichroism or infrared bands, but these quantities cannot be easily calculated from the protein configurations.³² Here, we use geometrical parameters related to the protein structure as order parameters. Figure 2(a–c) shows ensemble averages of the $rmsd$, a parameter that reports globally on the protein structure, and two single amino acid contacts that report on the local structure of the protein: the Trp-to-backbone




Figure 3

Superposition of representative structures belonging to the folded basins. Four representative structures are obtained by clustering the structures sampled at the lowest temperature (280 K) during the last 10 ns of the REMD simulation. The top most populated clusters correspond to structures at 0.075 nm (48%), 0.69 nm (7%), 0.15 nm (6%), 0.17 nm (5%), and 0.21 nm (4%) from structure 1 of the NMR ensemble. Cluster 2 (not shown) is assigned as an unfolded state. The plots show the superposition of the structures in the folded basin ($rmsd < 0.22$ nm). We used the Daura clustering method implemented in GROMACS.³³ (A) Tube representation of the backbone. (B) All nonhydrogen atoms. The structures representative of the top four clusters in the folded basins are labeled in cyan (first cluster), blue (third cluster), red (fourth cluster), and orange (fifth cluster). The main differences in the structures are in the 3–10 turn (Pro12-Ser13-Ser14-Gly15) region and the N- and C-termini.

hydrogen bond distance (to the Arg carbonyl) and the Asp-Arg ion pair distance distributions at various temperatures. The $rmsd$ distance distributions, shown in Figure 2(a), exhibit two large peaks for $rmsd$ below 0.22 nm; one at 0.06 nm and another at 0.18 nm. We assign these two peaks, characterized by $rmsd < 0.22$ nm, as the folded state consistent with our previous results using ff94.^{18,24} A peak at 0.52 nm builds up at higher temperatures, corresponds to the unfolded state. The smallest distance to the PDB structure 1 is 0.029 nm, the largest is 1.2 nm. An analysis of the 38 structures depos-

ited in the 1L2Y PDB file shows that these structures are at an average of 0.72 nm with a standard deviation of 0.022 nm.¹ The minimum distance between structures in the NMR ensemble is 0.0227 nm and the maximum is 0.150 nm. The NMR ensemble and the simulated distribution centered at 0.066 nm are very similar.

In addition to the $rmsd$ distance, we also monitor the presence of a hydrogen bond between the Ne atom of the Trp side chain and the backbone carbonyl oxygen of Asp. The distance distributions at various temperatures, shown in Figure 2(b), show a large peak centered at 0.3




Figure 4

Contour maps of the free energy of the Trp-cage as a function of the $rmsd$ and the radius of gyration (R_g) at 280, 320, and 350 K. The contour shows that the system occupies two main basins: one corresponding to the folded state (low $rmsd$ and compact states) and another at larger $rmsd > 0.4$ nm and less compact corresponding to the unfolded state. At even higher temperatures, the system becomes extended and only the large $rmsd$ and R_g basin is occupied. The folded basin for low $rmsd$ and R_g has two basins (substates) at low temperature. The legends use a logarithmic scale, with -10 corresponding to the highest probability and 0 to the lowest.




Figure 5

Fraction of replicas folded as a function of temperature.

(A) Comparison of the fraction of Trp-cage protein folded as a function of temperature for two different force fields, ff99SB (red and green curves) and ff94 (blue curve). The folding profiles for the ff99SB are practically indistinguishable for REMD simulation starting from completely unfolded states (red) and folded states (green). The ff99SB profiles cross the 50% fraction folded at $T = 321$ K in close agreement with experimental data ($T_{\text{exp}} = 317$ K). However, the calculated fraction of folded states is lower than measured values at lower T . (B) Fraction folded as a function of other order parameters that are representative of the folded Trp-cage folded structure. These parameters are $rmsd$ (red curve—folded is $rmsd < 0.22$ nm), fraction of a Trp-Ne to Asp backbone carbonyl hydrogen bond (blue curve, $d < 0.4$ nm), Arg-Asp ion pair (magenta curve, $d < 0.6$ nm), and number of alpha helical amino acids (divided by five to fit the same scale in the plot, green curve). An alpha helical amino acid is an amino acid with three consecutive ϕ angles in the alpha helical region ($\phi = -60^\circ \pm 30^\circ$ and $\phi = -47^\circ \pm 30^\circ$).³⁴ The maximum number of alpha helical segments observed is 10.

nm, a smaller peak centered at 0.5 nm, and a broad distribution for larger distances. The intensity of the 0.3-nm peak decreases drastically with increasing temperature, showing that we could also monitor this hydrogen bond to differentiate the folded and unfolded state. Another interaction that is associated with the stability of the

miniprotein is the Arg-Asp ion pair.^{1,18} The distance distributions for this ion pair [Fig. 2C] exhibit a similar behavior as the other distributions in Figure 2, in that the large peak at 0.43 nm decrease quickly in magnitude with increasing temperature. Figure 3 shows the superposition of structures belonging to the two folded substates. The main differences among the structures in the basin are in the turn (Pro12-Ser13-Ser14-Gly15) region and the N- and C- termini. The substates with larger $rmsd$ present in the folded basin also have a subpopulation in which the Trp-Ne H-bond (Supporting Information Fig. S9) is near 0.5 nm. In the thermodynamic analysis shown below, we assume that the folding/unfolding transition is two state. This is consistent with the thermodynamic analysis of Trp-cage⁹ but not with the fluorescence experiments,¹¹ which may be affected by the large reporter group attached to the protein.

Figure 4 shows the two-dimensional contours of conditional probabilities of $rmsd$ and the radius of gyration (R_g) of the molecule. These contours show that the system samples two basins, one describing the folded state (low R_g and $rmsd$) and another distribution centered at $rmsd \sim 0.6$ nm and $R_g \sim 0.8$ nm corresponding to the unfolded state. The unfolded state basin is not highly populated at low temperature but as the temperature increases above 315 K, it dominates the weight of the distribution. Within this order parameter representation, the folding/unfolding thermodynamics of the Trp-cage is two state like. An analysis of the distributions for $rmsd < 0.22$ nm shows that two substates compose this basin. Supporting Information Figure S10 shows time history of the $rmsd$ versus time for selected trajectories. Transitions between these two substates occur with a mean residence time in either state of 0.1 ns. Given the quick interconversion between these two states, we consider them to belong to one folded basin.

Equilibrium thermodynamics

Based on the decomposition of the free energy as a function of various order parameters, we can describe the thermodynamics of the system as a function of temperature, assuming the thermodynamics is two state. We find that a convenient parameter to characterize the population of folded and unfolded states is the $rmsd$ from the folded state. Figure 5(a) shows the fraction of configurations that are folded as a function of temperature. We see that the fraction of folded reaches a value near 75% at the lowest sampled temperature. The fraction folded reaches 50% (i.e., equilibrium between folded and unfolded) at $T = T_f = 321$ K (48°C) and decreases rapidly as a function of temperature. Here, T_f defines the folding temperature. A similar (almost indistinguishable curve) is obtained for simulations started from folded or unfolded states, as described in Figure 1(c). As a reference, we also show the fraction folded profile obtained

Table I
Thermodynamic Parameters for Trp-Cage

	Calculated	Experimental
T_0, P_0	298 K, 0.1 MPa	298 K, 0.1 MPa
$\Delta\alpha$ (kJ/mol)	0.027	Not available
$\Delta\beta$ (kJ/mol/MPa ²)	-0.025	Not available
ΔV_U (mL/mol)	-1.94	Not available
ΔS (kJ/mol/K)	0.053	0.16
ΔC_p (kJ/mol/K)	0.314	0.30 ± 0.1
ΔG_U (kJ/mol)	1.49	3.2
T_f (K)	321	317


Thermodynamic parameters describing the P - T stability diagram.³⁰

using the ff94 force field obtained previously.^{18,24} The ff94 force field yields a much more stable folded state with higher occupancy at low temperature (90%) and $T_f \sim 420$ K.

Similar thermodynamic profiles can be obtained using other order parameters. For example, monitoring the population of the hydrogen bond between the Trp-N ϵ to the Arg backbone carbonyl oxygen (labeled Trp-Hb) shows a very similar behavior as the *rmsd* order parameter. That is, the *rmsd* and hydrogen bond are formed at low temperature and the probability of forming the hydrogen bond decreases significantly as temperature is increased slightly. Figure 5(b) shows the fraction of the ensemble that forms the Trp-HB ($d < 0.40$ nm), the Arg-Asp ion pair ($d < 0.6$

nm), and the total number of alpha helical amino acids. The thermodynamic profile obtained with the Trp-HB is very similar to the one obtained with the *rmsd*. However, we see that the Arg-Asp ion pair and alpha helical formation persist at temperatures above T_f , and the thermodynamics profile obtained using this order parameter will be different than for the *rmsd* or Trp-HB parameters, described earlier. For this small protein, the ion pairs can be present even when the system is unfolded.

The free energy difference between the unfolded and the folded states are described by Eq. (1) with the coefficients described in Table I. The specific heat, ΔC_p , entropy, $\Delta S(T)$, and free energy, ΔG_0 , differences have been measured for Trp-cage by Streicher and Makhatadze⁹ and reported at $T = 298$ K and $P = 0.1$ MPa. The reported experimental values are also shown in Table I. We find that the fitted changes in specific heat and folding temperature are in close agreement with experiments. The calculated free energy difference is underestimated by <2 kJ/mol. Figure 6 shows a comparison of the calculated and the experimentally measured free energy, enthalpy and entropy as a function of temperature. We find a significant difference between the calculated and measured entropy and enthalpy changes on folding, which differ by close to a factor of two, but in such a way that there is a balance between the entropy and enthalpy that gives $\Delta G = 0$ at 321 K versus 317 K obtained in experiments.

**Figure 6**

Comparison of the computed (black solid line) and measured (red dotted line) (a) Free energy, (b) fraction folded, (c) entropy, and (d) enthalpy, as a function of temperature.




Figure 7

Stability of the Trp-cage protein as a function of temperature and pressure. (A) Average energy difference between unfolded and folded state ensembles calculated at various temperatures for a system at low pressure. (B) Average energy difference between unfolded and folded state ensembles calculated at various temperatures for a system at high pressure. The uncertainties are calculated over five, 100-ns long time segments. (C) Fraction of the ensemble folded as a function of temperature. The two curves show the results obtained from simulations at low and high average pressures. The solid lines are the best χ^2 fit of the thermodynamics of the system, following a fitting procedure described in the text. The parameters describing the changes in Gibbs free energy are shown in Table I. (D) Pressure-Temperature stability diagram for the Trp-cage protein. The contours for $\Delta G(T, P) = 0$ (contour closer to $P = 0$ and $T = 320$ K), -5 , -10 , and -15 kJ/mol are elliptical. The free energy profile shows small pressure dependence and the protein will require high pressures to unfold at low temperatures. The dotted lines represent the states sampled during the REMD simulations at low (blue) and high (red) densities. The black solid line represents the $\Delta V = 0$ isochore. The red solid line represents $\Delta S = 0$. All states above this curve decrease stability on cooling and have higher entropy in the folded state than in the unfolded state.

Figure 7 shows the fitting of the energy difference on unfolding at low (A) and high (B) densities. Notice that the fitted curve is almost linear over a broad range of temperatures. The large error bars at high temperatures are due to the lack of statistics for folded states at very temperatures ($T > 400$ K). Figure 7(C) shows the fraction of folded proteins as a function of temperature for the systems at low and high density (high average pressure). The curve shows that pressure destabilizes the protein. Pressure destabilization of the folded protein comes as a combination of many effects. On one hand, pressure will favor the state with lowest volume. There are three ways of reducing the volume of a protein: (i) by reducing the amount of free volume (cavities) in the folded state, (ii) by inserting water molecules into the hydrophobic core of the protein,³⁵ and (iii) by increasing the water density around nonpolar groups exposed to the solvent.^{24,36–38} The net equilibrium will be determined by the balance of these interactions in the folded and

unfolded states. From our free energy fits, the net volume change between the unfolded and folded state, ΔV_{U}^0 , is -2 mL/mol favoring the unfolded state. This small volume is consistent with the small hydrophobic core of the Trp-cage. Figure 7D shows the pressure-temperature (P - T) stability diagram for the Trp-cage. The $\Delta G(T, P) = 0$ curve is elliptical, suggesting that the protein can be denatured at low pressure by increasing the temperature (heat denaturation) or by increasing the pressure at constant low temperature (e.g., 298 K). We also see a region of $\Delta S(T, P) < 0$, which is a characteristic of cold denaturation at high pressures and low temperatures.

DISCUSSION AND CONCLUSIONS

We have used molecular dynamics simulations to describe the folding/unfolding equilibrium thermody-

namics for the Trp-cage miniprotein in water. The calculations exploit the use of enhanced sampling methods to accelerate the sampling of configurations while maintaining a Boltzmann distribution for the ensemble of configurations obtained. Comparison of the calculated folded state with the NMR structure shows that the ensemble of structures obtained are as close as 0.029 nm and that they share all secondary structural features and tertiary contacts characteristic of the Trp-cage (e.g., Arg-Asp ion pair, Trp-to-backbone hydrogen bond). The R_g of the unfolded state is slightly larger than for the folded state, but only at very high temperatures are noncollapsed states sampled. This is in agreement with photochemically induced NMR pulse-labeling experiments showing that non-native contacts between the Trp side chain protons and other side chains exist in the unfolded state.³⁹

The obtained configuration ensembles were analyzed within a two-state thermodynamics assumption to determine the pressure-temperature stability map of the system. This analysis describes an elliptical P - T diagram in which the protein can be unfolded via increases in pressure and/or temperature. The elliptical P - T diagrams have been observed for globular proteins.⁴⁰⁻⁴² For the Trp-cage, we find that the pressure effects are small. This may be a reflection of the fact that the protein does not have a large hydrophobic core, which when disrupted in the unfolded state, it helps to lower the overall volume of the system.³⁵ At 298 K and 0.1 MPa, the volume change, ΔV_0 , is negative and small (approximately -2 mL/mol). The MD simulations and modeling of the thermodynamics provides a model which is in excellent quantitative agreement with experiments.⁹ We find that the specific heat and free energy differences and folding temperature agree with experimental values to a degree unexpected for the model. Previous calculations of the thermodynamics using the ff94 force field produced similar results for the specific heat changes but gave the folding temperature at high values (~ 450 K).²⁴ The ff99SB force field used here gives much better results for the folding temperature. Comparison of MD simulations with NMR order parameter measurements have also shown that the ff99SB force field describes better, when compared to experiments, the dynamics of globular proteins.⁴³ However, studies of the helix-coil transition on model peptides have shown that the ff99SB force field fails to form thermodynamically stable helices at room temperature.⁴⁴ Given that the Trp-cage has a large helical segment, we could have expected that the ff99SB would not have folded. However, we find that for non-Ala-based sequences and in the presence of tertiary interactions, stable alpha helices are formed for the Trp-cage. The alpha helices are stable ~ 50 K above the folding T . The importance of tertiary contacts has been highlighted by Neidigh *et al.*¹ When the three C-terminal proresidues were removed from the Trp-cage construct, the N-terminal part of the peptide did show not any significant pro-

tection of the amide protons from H/D exchange.¹ We would like to point at another difference, structurally distinguishing the unfolded ensembles of the ff94 and ff99SB models: ff99SB shows an about a factor of four to five reduced helical content compared with ff94 at the lowest temperatures.²⁴ The residues involved in forming the N-terminal helix are significantly exploring beta, PPII and left-handed-helical parts of the backbone conformation space when in the unfolded state. We conjecture that this greater conformational diversity is a key factor for the dramatically slower convergence of the ff99SB REMD simulations.

A puzzling result we obtained is that the entropy and enthalpy changes on folding are at least a factor of two smaller than the measured values. Here, we estimate the entropies from the temperature dependence of the fitted free energies and not from the ensembles. Entropy and enthalpy changes on unfolding are typically an order of magnitude larger than the corresponding free energy differences,⁴⁵ and entropies cannot be calculated directly from average energies and free energies since the errors in average energy are as large as ΔG . Calculations based on minimalist models⁴⁶ and all-atom models²⁴ show, in agreement with experiments, that the entropy and enthalpies are approximately one order of magnitude larger than the free energies.⁴⁵ All atom calculations of enthalpies using molecular dynamics simulations of an RNA tetraloop hairpin also underestimated the enthalpy of the transition.⁴⁷ Best and Hummer optimized the dihedral angle energy terms of the ff99SB force field such that calculations could reproduce the known stability of alpha helices at a given temperature.⁴⁴ They found that, although they can reproduce the free energy at a given T , they could not get the correct temperature dependence. In the case of alpha helical peptides, it was also found that the entropy and enthalpies were underestimated by a factor of two. Best and Hummer have speculated that the smaller enthalpy and entropy changes may be a consequence of the simplicity of the energy model used in atomic simulations, in which hydrogen bonds are represented by Coulomb interactions and do not have a bonding character, and electronic polarizability are not explicitly included. We speculate that models that include directional hydrogen bonding terms, as they are used in structure prediction programs, may give a better description of the entropy and enthalpy differences.^{48,49} Interestingly, the underestimates in entropy and enthalpy are correlated such that the free energy differences are reasonably small and the folding transition temperature is close to the measured value. If we assume that the directional hydrogen bonding is responsible for the lower enthalpy, how do we justify the decrease in entropy? We can argue that nondirectional hydrogen bonds allow for more flexibility in the hydrogen bonds without much change in energy. If we use a directional hydrogen bond term, the energy will change on hydrogen bond bending

and the entropy of the folded state will be reduced. The reduction in entropy of the folded state will lead to a decrease in the stability of the folded state, unless a stronger binding energy is also used. Assuming that the directional hydrogen bonds have no effect in the entropy of the unfolded state, we could, qualitatively, account for the increase in energy and larger increase in entropy changes on unfolding. Directional hydrogen bonds, if used in the solvent model, will change the hydrophobic effect on folding by increasing solvent structure and, presumably, decreasing H-bonding between the protein and the solvent. These changes in the hydrophobic effect could also contribute to changes in entropy and enthalpy. A careful parameterization of the effect of directional hydrogen bonds will be needed such that the net changes in entropy and enthalpy cancel out in the free energy difference and give similar folding temperatures and free energy differences, as we obtained with the current force field. Based on the close agreement with experimental folding temperature, specific heat, and free energy differences, it could be argued that the ff99SB force field gives a better description of the thermodynamics than the ff94. However, there are differences in enthalpy and entropies, and as a consequence, the lack of sharp changes in the fraction of molecules in the folded state will need to be addressed in future parameterizations of empirical force fields. The effect of water model (e.g., TIP3P vs. TIP4P) on the folding/unfolding of the Trp-cage will be described in another publication.

ACKNOWLEDGMENTS

The authors thank Prof. George I. Makhatadze for helpful discussions.

REFERENCES

- Neidigh JW, Fesinmeyer RM, Andersen NH. Designing a 20-residue protein. *Nat Struct Biol* 2002;9:425–430.
- Qiu LL, Pabit SA, Roitberg AE, Hagen SJ. Smaller and faster: the 20-residue Trp-cage protein folds in 4 μ s. *J Am Chem Soc* 2002;124:12952–12953.
- Simmerling C, Strockbine B, Roitberg AE. All-atom structure prediction and folding simulations of a stable protein. *J Am Chem Soc* 2002;124:11258–11259.
- Hornak V, Abel R, Okur A, Strockbine B, Roitberg A, Simmerling C. Comparison of multiple amber force fields and development of improved protein backbone parameters. *Proteins* 2006;65:712–725.
- Zhan LX, Chen JZY, Liu WK. Computational study of the Trp-cage miniprotein based on the ECEPP/3 force field. *Proteins* 2007;66:436–443.
- Huang XH, Hagen M, Kim B, Friesner RA, Zhou RH, Berne BJ. Replica exchange with solute tempering: efficiency in large scale systems. *J Phys Chem B* 2007;111:5405–5410.
- Juraszek J, Bolhuis PG. Sampling the multiple folding mechanisms of Trp-cage in explicit solvent. *Proc Natl Acad Sci USA* 2006;103:15859–15864.
- Pitera JW, Swope W. Understanding folding and design: replica-exchange simulations of “Trp-cage” fly miniproteins. *Proc Natl Acad Sci USA* 2003;100:7587–7592.
- Streicher WW, Makhatadze GI. Unfolding thermodynamics of Trp-cage, a 20 residue miniprotein, studied by differential scanning calorimetry and circular dichroism spectroscopy. *Biochemistry* 2007;46:2876–2880.
- Nikiforovich GV, Andersen NH, Fesinmeyer RM, Frieden C. Possible locally driven folding pathways of TC5b, a 20-residue protein. *Proteins* 2003;52:292–302.
- Neuweiler H, Doose S, Sauer M. A microscopic view of miniprotein folding: enhanced folding efficiency through formation of an intermediate. *Proc Natl Acad Sci USA* 2005;102:16650–16655.
- Ahmed Z, Beta IA, Mikhonin AV, Asher SA. UV-resonance Raman thermal unfolding study of Trp-cage shows that it is not a simple two-state miniprotein. *J Am Chem Soc* 2005;127:10943–10950.
- Naduthambi D, Zondlo NJ. Stereoelectronic tuning of the structure and stability of the Trp-cage miniprotein. *J Am Chem Soc* 2006;128:12430–12431.
- Hudaky P, Straner P, Farkas V, Varadi G, Toth G, Perczel A. Cooperation between a salt bridge and the hydrophobic core triggers fold stabilization in a Trp-cage miniprotein. *Biochemistry* 2008;47:1007–1016.
- Bunagan MR, Yang X, Saven JG, Gai F. Ultrafast folding of a computationally designed Trp-cage mutant: Trp(2)-cage. *J Phys Chem B* 2006;110:3759–3763.
- Barua B, Lin JC, Williams VD, Kummeler P, Neidigh JW, Andersen NH. The Trp-cage: optimizing the stability of a globular miniprotein. *Protein Eng Des Sel* 2008;21:171–185.
- Seibert MM, Patriksson A, Hess B, van der Spoel D. Reproducible polypeptide folding and structure prediction using molecular dynamics simulations. *J Mol Biol* 2005;354:173–183.
- Paschek D, Nymeyer H, Garcia AE. Replica exchange simulation of reversible folding/unfolding of the Trp-cage miniprotein in explicit solvent: on the structure and possible role of internal water. *J Struct Biol* 2007;157:524–533.
- Patriksson A, Adams CM, Kjeldsen F, Zubarev RA, van der Spoel D. A direct comparison of protein structure in the gas and solution phase: the TRP-cage. *J Phys Chem B* 2007;111:13147–13150.
- Snow CD, Zagrovic B, Pande VS. The Trp cage: folding kinetics and unfolded state topology via molecular dynamics simulations. *J Am Chem Soc* 2002;124:14548–14549.
- Chowdhury S, Lee MC, Xiong GM, Duan Y. Ab initio folding simulation of the Trp-cage miniprotein approaches NMR resolution. *J Mol Biol* 2003;327:711–717.
- Zhou RH. Trp-cage: folding free energy landscape in explicit water. *Proc Natl Acad Sci USA* 2003;100:13280–13285.
- Juraszek J, Bolhuis PG. Rate constant and reaction coordinate of Trp-cage folding in explicit water. *Biophys J* 2008;95:4246–4257.
- Paschek D, Hempel S, Garcia AE. Computing the stability diagram Trp-cage miniprotein. *Proc Natl Acad Sci USA* 2008;105:17754–17759.
- Kannan S, Zacharias M. Folding simulations of Trp-cage miniprotein in explicit solvent using biasing potential replica-exchange molecular dynamics simulations. *Proteins* 2009;76:448–460.
- Cornell WD, Cieplak P, Bayley CI, Gould R, Merz KM, Ferguson DM, Spellmeyer DC, Fox T, Caldwell J, Kollman PA. A second generation force field for the simulation of proteins, nucleic acids, and organic molecules. *J Am Chem Soc* 1995;117:5179–5197.
- Jorgensen W, Chandrasekhar J, Madura J, Impey R, Klein M. Comparison of simple potential functions for simulating liquid water. *J Chem Phys* 1983;79:926–935.
- Cooke B, Schmidler SC. Preserving the Boltzmann ensemble in replica-exchange molecular dynamics. *J Chem Phys* 2008;129:164112 (1–17).
- Garcia AE, Hecce HD, Paschek D. Simulations of temperature and pressure unfolding of peptides and proteins with replica exchange molecular dynamics. *Annu Rep Comput Chem* 2006;2:83–95.
- Smeller L. Pressure-temperature phase diagrams of biomolecules. *Biochim Biophys Acta* 2002;1595:11–29.

31. Frenkel D, Smit B. Understanding molecular dynamics simulation: from algorithms to applications, 2nd ed. San Diego: Academic Press; 2002.
32. Gnanakaran S, Hochstrasser RM, Garcia AE. Nature of structural inhomogeneities on folding a helix and their influence on spectral measurements. *Proc Natl Acad Sci USA* 2004;101:9229–9234.
33. Daura X, Suter R, van Gunsteren WF. Validation of molecular simulation by comparison with experiment: rotational reorientation of tryptophan in water. *J Chem Phys* 1999;110:3049–3055.
34. Garcia AE, Sanbonmatsu KY. Alpha-helical stabilization by side chain shielding of backbone hydrogen bonds. *Proc Natl Acad Sci USA* 2002;99:2782–2787.
35. Hummer G, Garde S, Garcia AE, Paulaitis ME, Pratt LR. The pressure dependence of hydrophobic interactions is consistent with the observed pressure denaturation of proteins. *Proc Natl Acad Sci USA* 1998;95:1552–1555.
36. Ghosh T, Garcia AE, Garde S. Molecular dynamics simulations of pressure effects on hydrophobic interactions. *J Am Chem Soc* 2001;123:10997–11003.
37. Ghosh T, Garcia AE, Garde S. Enthalpy and entropy contributions to the pressure dependence of hydrophobic interactions. *J Chem Phys* 2002;116:2480–2486.
38. Ghosh T, Garcia AE, Garde S. Water-mediated three-particle interactions between hydrophobic solutes: size, pressure, and salt effects. *J Phys Chem B* 2003;107:612–617.
39. Mok KH, Kuhn LT, Goetz M, Day IJ, Lin JC, Andersen NH, Hore PJ. A pre-existing hydrophobic collapse in the unfolded state of an ultrafast folding protein. *Nature* 2007;447:106–109.
40. Brandts JF, Oliveira RJ, Westort C. Thermodynamics of protein denaturation effect of pressure on the denaturation of RNase A. *Biochemistry* 1970;9:1038–1047.
41. Panick G, Vidugiris GJA, Malessa R, Rapp G, Winter R, Royer CA. Exploring the temperature-pressure phase diagram of staphylococcal nuclease. *Biochemistry* 1999;38:4157–4164.
42. Herberhold H, Winter R. Temperature- and pressure-induced unfolding and refolding of ubiquitin: a static and kinetic Fourier transform infrared spectroscopy study. *Biochemistry* 2002;41:2396–2401.
43. Showalter SA, Bruschweiler R. Validation of molecular dynamics simulations of biomolecules using NMR spin relaxation as benchmarks: application to the AMBER99SB force field. *J Chem Theory Comput* 2007;3:961–975.
44. Best RB, Hummer G. Optimized molecular dynamics force fields applied to the helix-coil transition of polypeptides. *J Phys Chem B* 2009;113:9004–9015.
45. Makhatadze G, Privalov P. Energetics of protein structure. *Adv Protein Chem* 1995;47:307–425.
46. Nymeyer H, Garcia AE, Onuchic JN. Folding funnels and frustration in off-lattice minimalist protein landscapes. *Proc Natl Acad Sci USA* 1998;95:5921–5928.
47. Garcia AE, Paschek D. Simulation of the pressure and temperature folding/unfolding equilibrium of a small RNA hairpin. *J Am Chem Soc* 2008;130:815–817.
48. Kortemme T, Morozov AV, Baker D. An orientation-dependent hydrogen bonding potential improves prediction of specificity and structure for proteins and protein–protein complexes. *J Mol Biol* 2003;326:1239–1259.
49. Morozov AV, Kortemme T, Tsemekhman K, Baker D. Close agreement between the orientation dependence of hydrogen bonds observed in protein structures and quantum mechanical calculations. *Proc Natl Acad Sci USA* 2004;101:6946–6951.